



22-3-14

Complete Course™ Fertilizer

Homogeneous Small Particle Size

- Methylene urea for controlled nitrogen feeding on greens, tees, fairways, roughs and other turf areas.
- Provides a high rate of readily-available potassium for improved drought tolerance, greater disease resistance and improved wear tolerance.
- Small particle size for excellent distribution, even at reduced rates.

GENERAL INFORMATION

Particle Size:	SGN: 120-130 UI: 35+
Bulk Density:	47.0 lb/cu.ft.
Net Weight:	22.68 kg (50.0 lb)
Coverage:	1,022 m ² @ 0.49 kg N / 100 m ² (11,000 sq. ft. @ 1.0 lb N / 1,000 sq. ft.)

PRODUCT SPECIFICATIONS

ANALYSIS 22-3-14

N	22%	97% Urea, Ammoniacal, MDU, DMTU 3% WIN (1% HWSN) (2% HWIN)
P	3%	Monoammonium Phosphate
K	14%	Potassium Sulphate
S	10%	Ammonium Sulphate, Potassium Sulphate

22-3-14

Complete Course™ Fertilizer

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. EYE CONTACT WITH CONTENTS MAY CAUSE IRRITATION. FLUSH EYES WITH WATER AS SOON AS POSSIBLE.

GUARANTEED MINIMUM ANALYSIS:

Total Nitrogen (N).....	22%
4.0% total nitrogen from methylenediurea nitrogen	
0.9% total nitrogen from dimethylenetriurea nitrogen	
0.4% total nitrogen is water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	3%
Soluble potash (K ₂ O)	14%
Sulphur (S).....	10.0%

DIRECTIONS FOR USE:

Use anytime fine turf areas are actively growing where quick greening and elevated potassium fertilization is desired. Apply during the late summer or the fall months.

Apply at the **NORMAL RATE** of 0.49 kg N / 100 sq. metres (1.0 lb N / 1,000 sq. feet). Equivalent to 2.2 kg of product per 100 sq. metres or 4.5 lb of product per 1,000 sq. feet.

Apply to dry foliage. Water in after application to avoid possible footprinting.

When temperatures are above 30° C (85° F), apply at the **LIGHT RATE** of 0.24 kg N / 100 sq. metres (0.5 lb N / 1,000 sq. feet). Equivalent to 1.1 kg of product per 100 sq. metres (2.3 lb of product per 1,000 sq. ft.). Apply to dry foliage and water in following application.

During mid-summer, apply only at the **LIGHT RATE**. Apply to dry foliage. Water in following application. Mid-summer applications are not recommended for bentgrass and *Poa annua*.

STORAGE AND USE:

Store in a clean, dry place.
Reseal opened bag by folding top down and securing.
Use only clean, dry spreaders.
Do not leave unused product in spreaders.

FIRST AID:

IF SWALLOWED: Get medical attention or contact poison control centre.

IF IN EYES: Flush with plenty of water and get medical attention or contact poison control centre.

IF ON SKIN: Wash with soap and flush with plenty of water.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

SUGGESTED SPREADER SETTINGS

To provide proper distribution, calibrate spreader before application.

22.68 kg (50 lb) treats 2,042 sq. metres (22,000 sq. ft.)
at the **LIGHT RATE**

22.68 kg (50 lb) treats 1,022 sq. metres (11,000 sq. ft.)
at the **NORMAL RATE**

Spreader	Ground Speed	Width of Coverage	Light Rate	Normal Rate
ProTurf® Drop Type	4.8 km/hr (3 mph)	Overlap Wheels	4 ¼	5 ¼
ProTurf® Rotaries				
Accupro 2000 (Cone 6)	4.8 km/hr (3 mph)	3.4 m (11 feet)	H ½	J ½
SR 2000 (Cone 6)	4.8 km/hr (3 mph)	3.4 m (11 feet)	H ½	J ½
R-8, R-8A (Cone 6)	4.8 km/hr (3 mph)	3.0 m (10 feet)	H ½	J ½
Lely Models WTR, WFR, HR and 1250. PTO at 450 rpm.	7.25 km/hr (4.5 mph)	9.1 m (30 feet)	4 III	5 ¼ III
Vicon (03 Series- Standard Spout)	7.25 km/hr (4.5 mph)	8.5 m (28 feet)	17	25



ProTurf® is a registered trademark and Complete Course is a trademark, used under licence from The Scotts Company.
Nu-Gro Ltd., 10 Craig Street, Brantford, Ontario N3R 7J1

MAY 2006